

太空科學研究所 不分組 科目: 普通物理 共 / 頁 第 / 頁

1. A block of mass m is released on a wedge of mass M at a height h above the floor as in Fig.1. All surfaces are frictionless. Show that the speed of the wedge when the (25%) block hits the floor is given by

$$\sqrt{\frac{2m^2gh\cos^2\theta}{(M+m)(M+m\sin^2\theta)}}$$



- An ideal gas is taken around the cycle ABCA as shown in Fig.2. Find:
 - (a) the net work done by the gas in each cycle in terms of P_0 and V_0 ;
 - (b) the net heat input in each cycle.



(15%)

(15%)

Find the field due to an infinite flat sheet of charge with a uniform areal charge density $\sigma C/m^2$.

- (15%)
- A metal sphere of radius R has a charge Q. Find its potential energy.
- 5. A particle is subject to a constant force F along the direction of its motion. Starting with the expression $p = \gamma m_0 v$ and F = dp/dt, show that its acceleration is $dv/dt = F/\gamma^3 m_0$, where γ is the Lorentz factor.



An electron is accelerated from rest by a potential difference of 120 V. What (15%) is its de Broglie wavelength?

$$(m_e = 9.1 \times 10^{-31} \, kg, e = 1.6 \times 10^{-19} \, C, h = 6.6 \times 10^{-34} \, J \cdot s)$$