

※請在答案卡內作答

- 本測驗試題為複選題（答案可能有一個或多個），請選出所有正確或最適當的答案，並請用 2B 鉛筆作答於答案卡。
- 共二十題，每題完全答對得五分，答錯不倒扣。

Notation: In the following questions, underlined letters such as \underline{a} , \underline{b} , etc. denote column vectors of proper length; boldface letters such as \mathbf{A} , \mathbf{B} , etc. denote matrices of proper size; \mathbf{A}^T means the transpose of matrix \mathbf{A} . \mathbb{R} is the usual set of all real numbers. By $\mathbf{A} \in \mathbb{R}^{m \times n}$ we mean \mathbf{A} is an $m \times n$ real-valued matrix. $u(x)$ is unit-step function defined as $u(x) = 1$ if $x \geq 0$ and $u(x) = 0$ if $x < 0$; $*$ is the convolution operator; $\mathcal{L} : f(x) \mapsto F(s)$ and $\mathcal{L}^{-1} : F(s) \mapsto f(x)$ denote the unilateral Laplace and inverse Laplace transforms for $x \geq 0$, respectively.

- 一、 To solve a system of linear equations, we often use the augmented matrix form $[\mathbf{A} \ \underline{b}]$; then reduce \mathbf{A} to an upper triangular matrix by operating on the rows of \mathbf{A} and carry out the same operations on the vector \underline{b} . For instance, consider

$$\begin{cases} x + 2y + 2z = 1 \\ 4x + 8y + 9z = 3 \\ 3y + 2z = 1 \end{cases} \quad (1)$$

so we have

$$[\mathbf{A} \ \underline{b}] = \begin{bmatrix} 1 & 2 & 2 & 1 \\ 4 & 8 & 9 & 3 \\ 0 & 3 & 2 & 1 \end{bmatrix}.$$

Then we can apply an elimination matrix \mathbf{E} to the system such that $\mathbf{E}[\mathbf{A} \ \underline{b}] = [\mathbf{U} \ \underline{c}]$ and \mathbf{U} is an upper triangular matrix. Which of the following statements are true?

- (A) Left-multiply matrix \mathbf{A} by $\mathbf{E}_{21} = \begin{bmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ to subtract four times the first row of \mathbf{A} from the second row of \mathbf{A} .

- (B) Left-multiply matrix \mathbf{A} by $\mathbf{P}_{32} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ to exchange the second and the third columns of \mathbf{A} .

- (C) We can do both steps at once, i.e., set $\mathbf{E} = \begin{bmatrix} 1 & 0 & 0 \\ -4 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix}$ and $\mathbf{U} = \mathbf{E}\mathbf{A}$ is an upper triangular matrix.

- (D) The solution to the linear system (1) is $x = 1$, $y = 1$ and $z = -1$.

- (E) None of the above is true.

注意：背面有試題

參考用

類組：電機類 科目：工程數學 C(3005)

共 14 頁 第 2 頁

※請在答案卡內作答

二、 Suppose that matrix A can be factorized into the following form

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 5 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 & 5 \\ 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Which of the following statements are true?

- (A) $\text{rank}(A) = 2$
- (B) The dimension of the row-space of A equals 3.
- (C) The dimension of the column-space of A equals 2.
- (D) The dimension of the nullspace of A equals 1.
- (E) None of the above is true.

三、 Let $A, B \in \mathbb{R}^{4 \times 4}$ be matrices. Suppose that A and B have the same column space, but may not have the same columns. Which of the following statements are true?

- (A) Both matrices A and B have the same number of pivots.
- (B) Both matrices A and B have the same left nullspace and the same nullspace.
- (C) If A is invertible, so is B .
- (D) The row spaces of matrices A and B are orthogonal to each other under the usual Euclidean inner product.
- (E) None of the above is true.

注意：背面有試題

參考用

類組：電機類 科目：工程數學 C(3005)

共 14 頁 第 3 頁

※請在答案卡內作答

四、 Consider the matrix equation $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 1 & 1 \\ 1 & -1 \\ 2 & 0 \end{bmatrix} \quad \text{and } \mathbf{b} = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}$$

Let P be an orthogonal projection matrix from \mathbb{R}^3 onto the column space of A . Which of the following statements are true?

- (A) The matrix A is rectangular. It has no multiplicative left-inverse.
- (B) The solution to $A\hat{\mathbf{x}} = P\mathbf{b}$ is $\hat{\mathbf{x}} = [1 \ 1]^T$.
- (C) The projection vector $P\mathbf{b} = \frac{1}{3}[2 \ 3]^T$.
- (D) $P = A(A^T A)^{-1}A^T$; P is symmetric and satisfies $P^2 = P$.
- (E) None of the above is true.

五、 For the 4×4 matrix

$$A = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 2 & 1 & 3 \\ 1 & 3 & 1 & 9 \end{bmatrix}$$

which of the following statements are true?

- (A) $\det(A) = 3$
- (B) A is invertible
- (C) $\det(A^{-1}) = 3$
- (D) The (1,2)th entry of A^{-1} equals 3.
- (E) None of the above is true.

注意：背面有試題

參考用

※請在答案卡內作答

- 六、 What are the coordinate vectors for $f(x) = 5x^2 + 2x - 3$, with ordered bases $\mathcal{B}_1 = \{x^2, x, 1\}$ and $\mathcal{B}_2 = \{x^2 - x + 1, 3x^2 + 1, 2x^2 + x - 2\}$, respectively?

$$(A) [f(x)]_{\mathcal{B}_1} = \begin{bmatrix} -3 \\ 2 \\ 5 \end{bmatrix}, [f(x)]_{\mathcal{B}_2} = \begin{bmatrix} 0 \\ -\frac{13}{3} \\ \frac{5}{3} \end{bmatrix}$$

$$(B) [f(x)]_{\mathcal{B}_1} = \begin{bmatrix} 0 \\ -\frac{13}{3} \\ \frac{5}{3} \end{bmatrix}, [f(x)]_{\mathcal{B}_2} = \begin{bmatrix} 5 \\ 2 \\ -3 \end{bmatrix}$$

$$(C) [f(x)]_{\mathcal{B}_1} = \begin{bmatrix} \frac{2}{3} \\ -\frac{13}{3} \\ 0 \end{bmatrix}, [f(x)]_{\mathcal{B}_2} = \begin{bmatrix} 5 \\ 2 \\ -3 \end{bmatrix}$$

$$(D) [f(x)]_{\mathcal{B}_1} = \begin{bmatrix} \frac{2}{3} \\ -\frac{13}{3} \\ 0 \end{bmatrix}, [f(x)]_{\mathcal{B}_2} = \begin{bmatrix} 2 \\ 5 \\ -3 \end{bmatrix}$$

- (E) None of the above is true.

- 七、 Given a non-zero matrix $A \in \mathbb{R}^{n \times n}$, which of the following statements are true?

- (A) The nullspace of A equals the left nullspace of A .
 (B) The eigenvectors of A are linearly independent.
 (C) Part of the eigen-space can equal the left nullspace of A .
 (D) The dimension of the eigen-space of A equals n .
 (E) None of the above is true.

注意：背面有試題

參考用

類組：電機類 科目：工程數學 C(3005)

共 14 頁 第 5 頁

※請在答案卡內作答

八、Let T be a linear operator on \mathbb{R}^3 defined as

$$T(\underline{x}) = \begin{bmatrix} 2x_2 + x_3 \\ x_1 - 4x_2 \\ 3x_1 \end{bmatrix}, \text{ with } \underline{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}.$$

What is the standard matrix, $[T]_{\mathcal{B}}$, associated to T with respect to the ordered basis

$$\mathcal{B} = \left\{ \underline{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \underline{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \underline{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\},$$

and $[T(\underline{x})]_{\mathcal{B}}$?

$$(A) [T]_{\mathcal{B}} = \begin{bmatrix} 0 & 2 & 3 \\ 1 & -4 & 0 \\ 3 & 0 & 0 \end{bmatrix}, [T(\underline{x})]_{\mathcal{B}} = \begin{bmatrix} x_3 \\ x_2 - x_3 \\ x_1 - x_2 \end{bmatrix}.$$

$$(B) [T]_{\mathcal{B}} = \begin{bmatrix} 3 & 3 & 3 \\ -6 & -6 & -2 \\ 6 & 5 & -1 \end{bmatrix}, [T(\underline{x})]_{\mathcal{B}} = \begin{bmatrix} 3x_1 \\ -2x_1 - 4x_2 \\ -x_1 + 6x_2 + x_3 \end{bmatrix}.$$

$$(C) [T]_{\mathcal{B}} = \begin{bmatrix} 3 & 3 & 3 \\ -6 & -6 & -2 \\ 6 & 5 & -1 \end{bmatrix}, [T(\underline{x})]_{\mathcal{B}} = \begin{bmatrix} x_3 \\ x_2 - x_3 \\ x_1 - x_2 \end{bmatrix}.$$

$$(D) [T]_{\mathcal{B}} = \begin{bmatrix} 0 & 2 & 3 \\ 1 & -4 & 0 \\ 3 & 0 & 0 \end{bmatrix}, [T(\underline{x})]_{\mathcal{B}} = \begin{bmatrix} 3x_1 \\ -2x_1 - 4x_2 \\ -x_1 + 6x_2 + x_3 \end{bmatrix}.$$

(E) None of the above is true.

注意：背面有試題

參考用

※請在答案卡內作答

九、 Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ be a non-zero, symmetric matrix. Suppose the eigen-decomposition of $\mathbf{A} = \mathbf{Q}\mathbf{\Lambda}\mathbf{Q}^T$, where $\mathbf{\Lambda}$ is a diagonal matrix and $\mathbf{Q} = \begin{bmatrix} \mathbf{Q}_1 & \mathbf{Q}_2 \end{bmatrix}$ is an orthonormal matrix, with $\mathbf{Q}_1 \in \mathbb{R}^{n \times r}$ and $\mathbf{Q}_2 \in \mathbb{R}^{n \times (n-r)}$, $r = \text{rank}(\mathbf{A})$. Which of the following statements are true?

- (A) $\text{span}(\mathbf{x} - \mathbf{Q}_1\mathbf{Q}_1^T\mathbf{x}) = \text{span}(\mathbf{Q}_2\mathbf{Q}_2^T\mathbf{x})$ for any given vector $\mathbf{x} \in \mathbb{R}^n$.
- (B) $\mathbf{A}\mathbf{Q}_2$ is not an all-zero matrix.
- (C) $\mathbf{Q}_1^T\mathbf{Q}_2$ is an all-zero matrix.
- (D) $\mathbf{Q}_2^T\mathbf{Q}_2$ is an all-zero matrix.
- (E) None of the above is true.

十、 Which of the following statements are true?

- (A) A real matrix may be positive definite without being symmetric with respect to the real Euclidean inner product.
- (B) A positive semidefinite matrix cannot have 0 as an eigenvalue.
- (C) $\det : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is a linear transformation.
- (D) Let \mathbf{A} and \mathbf{B} be square matrices; then $\det(\mathbf{A}^T\mathbf{B}) = \det(\mathbf{A})\det(\mathbf{B})$.
- (E) None of the above is true.

注意：背面有試題

參考用

※請在答案卡內作答

十一、Consider the following non-homogeneous system:

$$\begin{cases} y_1'(x) = 4y_1(x) + 2y_2(x) - 15xe^{-2x} \\ y_2'(x) = 3y_1(x) - y_2(x) - 4xe^{-2x} \end{cases}$$

satisfying $y_1(0) = 7$ and $y_2(0) = 3$. Which of the following statements are true?

- (A) $y_1(x) = (6 + 2x - 7x^2)e^{-2x} + e^{5x}$ and $y_2(x) = (-4 + 21x^2)e^{-2x} + 7e^{5x}$.
- (B) $y_1(x) = \frac{14+13x^2}{14}e^{-2x} + \frac{42+4x}{7}e^{5x}$ and $y_2(x) = \frac{28+5x}{14}e^{-2x} + \frac{7+10x+6x^2}{7}e^{5x}$.
- (C) $y_1(1) = \frac{27}{14}e^{-2} + \frac{46}{7}e^5$ and $y_2(1) = \frac{31}{14}e^{-2} + \frac{23}{7}e^5$.
- (D) $y_1(-1) = -3e^2 + e^{-5}$ and $y_2(-1) = 17e^2 + 7e^{-5}$.
- (E) None of the above is true.

十二、Consider the differential equation $(2xy(x) - x^2 - y^2(x) - 1)dx + dy(x) = 0$. Which of the following statements are true?

- (A) The differential equation is exact.
- (B) $x + y(x) = 0$ is an implicit solution.
- (C) If $y(0) = \frac{1}{2}$, then $y(1) = 2$.
- (D) If $y(0) = 1$, then $y(2) = 1$.
- (E) None of the above is true.

注意：背面有試題

參考用

※請在答案卡內作答

十三、 Consider the differential equation $xdy(x) - 2y(x)[2x^2 - \ln y(x)]dx = 0$. Which of the following statements are true?

- (A) The differential equation is linear.
- (B) $y(x) = e^{x^2}$ is a particular solution.
- (C) If $y(1) = 1$, then $y(2) = \frac{15}{4}$.
- (D) If $y(1) = 1$, then $y'(1) = 4$.
- (E) None of the above is true.

十四、 Consider the differential equation $(1 - x^2)y''(x) - 2xy'(x) + 2y(x) = 0$. Which of the following statements are true?

- (A) $y(x) = \ln \frac{1+x}{1-x}$ is a particular solution.
- (B) $y(x) = x$ is a particular solution.
- (C) If $y(0) = 1$ and $y'(0) = 1$, then $y(\frac{1}{2}) = \frac{3}{2} - \frac{\ln 3}{4}$.
- (D) If $y(0) = 2$ and $y'(0) = 0$, then $y(\frac{1}{2}) = 2 + \ln 3$.
- (E) None of the above is true.

注意：背面有試題

參考用

※請在答案卡內作答

十五、 Consider the following differential equation:

$$(x^3+1)y^{(3)}(x)+(3-x^2)y''(x)+(4+2x)y'(x)+10y(x) = 4+12x+(4x-8x^3)\cos(2x)+(4x^2-2)\sin(2x)$$

satisfying $y(\pi) = \pi$, $y'(\pi) = 3$ and $y''(\pi) = 0$. Which of the following statements are true?

- (A) $y(0) = 0$.
- (B) $y(\frac{\pi}{2}) = \frac{\pi}{2}$.
- (C) $y'(\frac{\pi}{4}) = 1$.
- (D) $y'(1) = \pi$.
- (E) None of the above are true.

注意：背面有試題

參考用

※請在答案卡內作答

十六、 Consider the system of differential equations $\underline{y}'(x) = \mathbf{A}\underline{y}(x) + \underline{f}(x)$, where

$$\mathbf{A} = \begin{bmatrix} 2 & -4 \\ 1 & -2 \end{bmatrix}, \underline{y}(0) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \text{ and } \underline{f}(x) = \begin{bmatrix} x^2 \\ x \end{bmatrix}.$$

Which of the following statements are true?

- (A) $e^{\mathbf{A}x}\underline{y}(0) = \begin{bmatrix} 1 - 3x \\ 1 - x \end{bmatrix}$
- (B) $\mathcal{L}\{\underline{y}(x)\} = \frac{1}{s^3} \begin{bmatrix} s^4 - 2s^3 + 2s + 4 \\ s^4 - s^3 + s^2 - 2s + 2 \end{bmatrix}$
- (C) $\underline{y}(1) = \left[-\frac{7}{6} \quad \frac{1}{4}\right]^T$
- (D) $[2 \ 1]^T$ is a generalized eigenvector for matrix \mathbf{A} based on eigenvector $[1 \ 0]^T$.
- (E) None of the above is true.

十七、 Let $g(x)$ be a real-valued function given as below

$$g(x) = \begin{cases} e^{-|x|}, & \text{if } x \in (-\pi, \pi), \\ 0, & \text{otherwise} \end{cases}$$

which has a Fourier series (F.S.) representation for $x \in (-\pi, \pi)$ as follows

$$g(x) \stackrel{\text{F.S.}}{=} g_0 + \sum_{n \geq 1} [a_n \cos(nx) + b_n \sin(nx)]$$

Which of the following statements are true?

注意：背面有試題

參考用

※請在答案卡內作答

- (A) $g_0 = \frac{1-e^{-\pi}}{\pi}$
 (B) $a_1 = \frac{1-e^{-\pi}}{\pi}$
 (C) $b_1 = 0$
 (D) $2g_0^2 + \sum_{n \geq 1} (a_n^2 + b_n^2) = \frac{1-e^{-2\pi}}{\pi}$
 (E) None of the above is true.

十八、Continued from Question 十七, solve for $y(x)$ the following differential equation

$$y''(x) - 9y(x) = \sum_{n=-\infty}^{\infty} g(x - 2n\pi).$$

for all $x \in \mathbb{R}$ and $x \neq \pm\pi, \pm3\pi, \pm5\pi, \dots$. With $y(0) = y'(0) = 0$, it is already known that the solution $y(x)$ takes the following form

$$y(x) \stackrel{\text{F.S.}}{=} y_0 + \sum_{n \geq 1} [c_n \cos(nx) + d_n \sin(nx)]$$

Which of the following statements are true?

- (A) $y_0 = -\frac{1-e^{-\pi}}{9\pi}$
 (B) $c_n < 0$ for all $n \geq 1$
 (C) For $n \geq 1$, we have $\frac{|a_n|}{|c_n|} = n^2 + 9$.
 (D) With the same initial conditions, the solution $y(x)$ can be alternatively represented as

$$y(x) = f(x) * \left(\sum_{n=-\infty}^{\infty} \delta(x - 2n\pi) \right)$$

注意：背面有試題

參考用

類組：電機類 科目：工程數學 C(3005)

共 14 頁 第 12 頁

※請在答案卡內作答

where

$$f(x) = \left[\frac{1}{8} \cosh(3x) - \frac{2u(x) - 1}{24} \sinh(3x) - \frac{1}{8} e^{-|x|} \right] (u(x + \pi) - u(x - \pi))$$

(E) None of the above is true.

注意：背面有試題

參考用

類組：電機類 科目：工程數學 C(3005)

共 14 頁 第 13 頁

※請在答案卡內作答

十九、Solve the following boundary value problem for the real-valued bi-variate function $f(x, y)$ for $0 \leq x \leq 2\pi$ and $y \geq 0$ satisfying

$$\frac{\partial^2 f(x, y)}{\partial x^2} = 4 \frac{\partial^2 f(x, y)}{\partial y^2}$$

$$f(0, y) = f(2\pi, y) = f(x, 0) = 0, \quad \left. \frac{\partial f(x, y)}{\partial y} \right|_{y=0} = [\sin(x)]^3.$$

Which of the following statements are true?

- (A) $f\left(\frac{\pi}{2}, \pi\right) = \frac{4}{3}$.
- (B) $f\left(\frac{\pi}{2}, \frac{\pi}{2}\right) = -\frac{5\sqrt{2}}{6}$.
- (C) $\left. \frac{\partial f(x, y)}{\partial y} \right|_{x=y=\frac{\pi}{2}} = \frac{\sqrt{2}}{4}$
- (D) $\left. \frac{\partial^2 f(x, y)}{\partial y^2} \right|_{x=y=\frac{\pi}{2}} = \frac{3\sqrt{2}}{8}$
- (E) None of the above is true.

注意：背面有試題

參考用

類組：電機類 科目：工程數學 C(3005)

共 14 頁 第 14 頁

※請在答案卡內作答

二十、Continued from Question 十九, if we change the boundary value problem to

$$\frac{\partial^2 f(x, y)}{\partial x^2} + 4 \frac{\partial^2 f(x, y)}{\partial y^2} = 0$$

while keeping all the boundary- and initial conditions the same for $f(x, y)$, which of the following statements are true?

- (A) $\left. \frac{\partial f(x, y)}{\partial y} \right|_{x=\frac{\pi}{4}, y=0} = \frac{\sqrt{2}}{2}$
- (B) $\left. \frac{\partial f(x, y)}{\partial y} \right|_{x=\frac{\pi}{2}, y=0} = 1$
- (C) $\left. \frac{\partial^2 f(x, y)}{\partial x^2} \right|_{y=0} = 1$
- (D) $\lim_{y \rightarrow \infty} \frac{\ln(|f(\frac{3\pi}{4}, y)|)}{y} = \frac{3}{2}$.
- (E) None of the above is true.

注意：背面有試題

參考用