

系所別: 工業管理研究所甲、乙組 科目:


統計學

1. (10%) Explain the relationship between *Poisson* distribution and *exponential* distribution.
2. (10%) Explain the relationship between *Gamma* distribution, *m-Erlang* distribution, and *exponential* distribution.
3. (10%) Explain the relationship between *exponential* distribution and *Weibull* distribution.
4. (10%) Explain the relationship between *Bernoulli* distribution and *binomial* distribution.
5. (20%) Is the type of beverage ordered with lunch at a restaurant independent of the age of the consumer? A random pool of 259 lunch customers is taken, resulting in the following contingency table of observed values. Use $\alpha = 0.01$ to determine whether the two variables are independent. (Hint: $\chi^2 = \sum \sum \frac{(f_o - f_e)^2}{f_e}$)

參考用

Preferred Beverage

	Preferred Beverage			
	Coffee/Tea	Soft Drink	Other (Milk, etc.)	
21-34	26	45	18	89
35-55	41	40	20	101
>55	24	13	32	69
	91	98	70	259



Example: df (Number of degrees of freedom) = 5, the tail above $\chi^2 = 9.23635$ represents 0.10 or 10% of the area under the curve.

AREA IN UPPER TAIL

Degrees of Freedom	.995	.99	.975	.95	.90	.10	.05	.025	.01	.005
1	392.702×10^{-10}	187.083×10^{-9}	982.069×10^{-8}	993.214×10^{-8}	.0167908	2.70554	3.84146	5.02389	6.63490	7.87944
2	.0100251	.0201007	.0506356	.102587	.210730	4.60517	5.99147	7.37776	9.21034	10.5966
3	.0717212	.14832	.215795	.351846	.584375	6.25139	7.81473	9.34840	11.3449	12.8381
4	.206990	.297110	.484419	.710721	1.063623	7.77944	9.48773	11.1433	13.2767	14.8602
5	.411740	.554300	.831211	1.145476	1.61031	9.23635	11.0705	12.8325	15.0863	16.7496
6	.675727	.872085	1.237347	1.63539	2.20413	10.6446	12.5916	14.4494	16.8119	18.5476
7	.989265	1.239043	1.68987	2.16735	2.83311	12.0170	14.0671	16.0128	18.4753	20.2777
8	1.344419	1.646482	2.17973	2.73264	3.48954	13.3616	15.5073	17.5346	20.0902	21.9550
9	1.734926	2.087912	2.70039	3.32511	4.16816	14.6837	16.9190	19.0228	21.6660	23.5893

注意: 背面有試題


系所別: 工業管理研究所甲、乙組 科目:

統計學

6. (20%) If 10% of a population of parts is defective, what is the probability of randomly selecting 80 parts and finding that 12 or more parts are defective?
7. (20%) A sample of 87 professional working women showed that the average amount paid annually into a private pension fund per person was \$3343, with a sample standard deviation of \$1226. A Sample of 76 professional working men showed that the average amount paid annually into a private pension fund per person was \$5568, with a sample standard deviation of \$1716. A women's activist group wants to "prove" that women do not pay as much annually as men into private pension funds. If they use $\alpha = 0.001$ and these sample data, will they be able to reject a null hypothesis that women annually pay the same as or more than men into private pension funds? Use the step-by-step hypothesis-testing process.

TABLE A.1 Cumulative Probabilities of the Standard Normal Distribution

Entry is area A under the standard normal curve from $-\infty$ to $z(A)$



z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9971	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998



Selected Percentiles	
Cumulative probability A:	z(A):
.90	1.282
.95	1.645
.975	1.960
.98	2.054
.99	2.326
.995	2.576
.999	3.090