

科目：物理化學(1004)校系所組：中央大學化學學系交通大學應用化學系 (甲組)清華大學化學系清華大學材料科學工程學系 (丙組)**1. (20 %, 5 points for each)**

- (1-1) Express the definition of the Gibbs free energy.
(1-2) Under what condition, an endothermic reaction can proceed spontaneously?
(1-3) Under what condition, an exothermic reaction cannot proceed spontaneously?
(1-4) What are the second and third laws of thermodynamic?

2. (10 %, 5 points for each)

A sample of potassium ($K_{(s)}$, atomic weight = 23) of mass 2.140 g undergoes combustion in a constant-volume calorimeter with a calorimeter constant $1849 \text{ J} \cdot \text{K}^{-1}$ at room temperature. After the reaction completes, the temperature of the calorimeter and the inner water bath that contains 1450 g of water increases by 2.62 K. (1 cal = 4.184 J)

- (2-1) Calculate ΔU_f° for K_2O .
(2-2) Calculate ΔH_f° for K_2O .

3. (20 %, 5 points for each)

Consider a reaction: $2A \rightarrow 3B$. Given that the standard molar enthalpies of the reactant at T_1 (=273 K) and T_2 (=373 K) are $-65 \text{ kJ} \cdot \text{mol}^{-1}$ and $435 \text{ kJ} \cdot \text{mol}^{-1}$, respectively. Also, the standard molar enthalpy of the product at T_1 and its molar heat capacity are $-110 \text{ kJ} \cdot \text{mol}^{-1}$ and $22 \text{ kJ} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$, respectively. Assume that the enthalpies vary linearly with temperature.

- (3-1) Calculate the molar heat capacity of the reactant.
(3-2) Calculate the total enthalpy of the product at T_2 .
(3-3) Derive and determine if the reaction at T_1 and T_2 is endothermic or exothermic.
(3-4) Sketch in a plot showing how the total enthalpy of the reactants (2A) and that of the products (3B) vary with temperature.

注：背面有試題

科目：物理化學(1004)

校系所組：中央大學化學學系

交通大學應用化學系(甲組)

清華大學化學系

清華大學材料科學工程學系(丙組)

4. (26 %)

The state of an electron in a hydrogen atom is specified by three quantum numbers n , l , and m as

$$\psi_{nlm}(r, \theta, \phi) = R_{nl}(r)Y_{lm}(\theta, \phi)$$

where $R_{nl}(r)$ and $Y_{lm}(\theta, \phi)$ are the radial and angular parts, respectively. Its energy is

$$E_n = \frac{hcR_H}{n^2}$$

where h is Planck's constant, c is the speed of light, and R_H is the Rydberg constant for hydrogen ($R_H = 1.097 \times 10^5 \text{ cm}^{-1}$).

- (4-1) What is the quantum number n called? (4 points)
 (4-2) What is the degeneracy of the $n = 2$ level? (4 points)
 (4-3) For $l = 0, 1, \text{ and } 2$, the corresponding subshells are denoted as $s, p, \text{ and } d$, respectively. Which of the following transition(s) is not observed, in principle, in emission spectra? (6 points)
 A) $5p \rightarrow 1s$ B) $4d \rightarrow 3s$ C) $4s \rightarrow 2p$ D) $5p \rightarrow 3d$ E) $3s \rightarrow 4p$
 (4-4) Determine the longest possible wavelength and the shortest possible wavelength (the series limit) for lines in the Balmer series (series where the final state is the $n = 2$ level) of the spectrum of atomic hydrogen. Give your answer in units of nm. (12 points)

5. (24 %)

Consider a system of N distinguishable particles having only two energy levels 0 and ε ($\varepsilon > 0$): Suppose this system is in thermal equilibrium at temperature T .

- (5-1) Find the canonical partition function Q . Use the Boltzmann constant k . (8 points)
 (5-2) Find the internal energy U of the system. (8 points)
 (5-3) Find the isochoric heat capacity C_V of the system. (8 points)